

- lism in severely burned patients. *Metabolism* 1979; 28: 1031-1039
4. Wolfe RR, O'Donnell TF Jr, Stone MD, Richmand DA, Burke JF: Investigation of factors determining the optimal glucose infusion rate in total parenteral nutrition. *Metabolism* 1980; 29: 892-900
 5. Shaw JH, Wolfe RR: Response to glucose and lipid infusions in sepsis: a kinetic analysis. *Metabolism* 1985; 34: 442-449
 6. Wolfe RR, Shaw JHF: Glucose and FFA kinetics in sepsis: role of glucagon and sympathetic nervous system activity. *Am J Physiol* 1985; 248: E236-E243
 7. Shaw JH, Wolfe RR: Fatty acid and glycerol kinetics in septic patients and in patients with gastrointestinal cancer. The response to glucose infusion and parenteral feeding. *Ann Surg* 1987; 205: 368-376
 8. Shaw JHF, Humberstone DA: Cancer: a metabolic parasite. *Br J Surg* 1988; 75: 1262
 9. Sakurai Y, Klein S: Metabolic alteration in patients with cancer-nutritional implications. *Surg Today* 1998; 28: 247-257
 10. Kinney JM: Metabolic responses to injuries. In: Winters RW, Green HL, eds, *Nutritional Support of the Seriously Ill Patient*, New York, Academic Press, 1983; 5-12
 11. Wolfe RR, Goodenough RD, Burke JF, Wolfe MH: Response of protein and urea kinetics in burn patients to different levels of protein intake. *Ann Surg* 1983; 197: 163-171
 12. Shangraw RE, Stuart CA, Prince MJ, Peters EJ, Wolfe RR: Insulin responsiveness of protein metabolism *in vivo* following bedrest in humans. *Am J Physiol* 1988; 255: E548-E558
 13. Shangraw RE, Jahoor F, Miyoshi H, Neff WA, Stuart CA, Herndon DN, Wolfe RR: Differentiation between septic and postburn insulin resistance. *Metabolism* 1989; 38: 983-989
 14. Jahoor F, Shangraw RE, Miyoshi H, Wallfish H, Herndon DN, Wolfe RR: Role of insulin and glucose oxidation in mediating the protein catabolism of burns and sepsis. *Am J Physiol* 1989; 257: E323-E331
 15. Wolfe RR, Jahoor F, Herndon DN, Miyoshi H: Isotopic evaluation of the metabolism of pyruvate and related substrates in normal adult volunteers and severely burned children: effect of dichloroacetate and glucose infusion. *Surgery* 1991; 110: 54-67
 16. Wolfe RR, Jahoor F, Hartl WH: Protein and amino acid metabolism after injury. *Diabetes Metab Rev* 1989; 5: 149-164
 17. Bilmazes C, Kien CL, Rohrbach DK, Uauy R, Burke JF, Munro HN, Young VR: Quantitative contribution by skeletal muscle to elevated rates of whole-body protein breakdown in burned children as measured by N tau-methylhistidine output. *Metabolism* 1978; 27: 671-676
 18. Long CL, Birkhahn RG, Geiger JW, Betts JE, Schiller WR, Blakemore WS: Urinary excretion of 3-methylhistidine: an assessment of muscle protein catabolism in adult normal subjects and during malnutrition, sepsis, and skeletal trauma. *Metabolism* 1981; 30: 765-776
 19. Jeevanandam M, Horowitz GD, Lowry SF, Brennan MF: Cancer cachexia and protein metabolism. *Lancet* 1984; 1: 1423-1426
 20. Burt ME, Stein TP, Brennan MF: A controlled, randomized trial evaluating the effects of enteral and parenteral nutrition on protein metabolism in cancer-bearing man. *J Surg Res* 1983; 34: 303-314
 21. Shaw JH, Wolfe RR: Whole-body protein kinetics in patients with early and advanced gastrointestinal cancer: the response to glucose infusion and total parenteral nutrition. *Surgery* 1988; 103: 148-155
 22. Shaw JH, Wolfe RR: Metabolic intervention in surgical patients. An assessment of the effect of somatostatin, ranitidine, naloxone, diclofenac, dipyridamole, or salbutamol infusion on energy and protein kinetics in surgical patients using stable and radioisotopes. *Ann Surg* 1988; 207: 274-282
 23. Wolfe RR: Counterregulatory hormones and human protein metabolism. In: Nair KS, ed, *Protein Metabolism in Diabetes Mellitus*, London, Smith Gordon & Company, 1992; 207-213
 24. Moldawer LL, Svaninger G, Gelin J, Lundholm KG: Interleukin 1 and tumor necrosis factor do not regulate protein balance in skeletal muscle. *Am J Physiol* 1987; 253: C766-C773
 25. Scrimshaw NS, Hussein MA, Murray E, Rand WM, Young VR: Protein requirements of man: variations in obligatory urinary and fecal nitrogen losses in young men. *J Nutr* 1972; 102: 1595-1604
 26. Hegsted DM: Assessment of nitrogen requirements. *Am J Clin Nutr* 1978; 31: 1669-1677
 27. Picou D, Taylor-Roberts T: The measurement of total protein synthesis and catabolism and nitrogen turnover in infants in different nutritional states and receiving different amounts of dietary protein. *Clin Sci* 1969; 36: 283-296
 28. Wolfe RR, ed: *Radioactive and Stable Isotope Tracers in Biomedicine: Principles and Practice of Kinetic Analysis*, New York, Wiley-Liss, 1992
 29. Shaw JH, Wolfe RR: Glucose and urea kinetics in patients with early and advanced gastrointestinal cancer: the response to glucose infusion, parenteral feeding, and surgical resection. *Surgery* 1987; 101: 181-191
 30. Carraro F, Kimbrough TD, Wolfe RR: Urea kinetics in humans at two levels of exercise intensity. *J Appl Physiol* 1993; 75: 1180-1185
 31. Rennie MJ: Muscle protein turnover and wasting due to injury and disease. *Br Med Bull* 1985; 41: 257-264
 32. Waterlow JC, Garlick PJ, Millward DJ, eds: *Protein Turnover in Mammalian Tissues and in the Whole Body*, Amsterdam, North-Holland, 1978; 225-300
 33. Garlick PJ, Wernerman J, McNurlan MA, Essen P, Lobleby GE, Milne E, Calder GA, Vinnars E: Measurement of the rate of protein synthesis in muscle of postabsorptive young men by injection of a 'flooding dose' of [^{13}C] leucine. *Clin Sci* 1989; 77: 329-336
 34. Jahoor F, Zhang XJ, Baba H, Sakurai Y, Wolfe RR: Comparison of constant infusion and flooding dose techniques to measure muscle protein synthesis rate in dogs. *J Nutr* 1992; 122: 878-887
 35. Garlick PJ, Grant I: Amino acid infusion increases the sensitivity of muscle protein synthesis *in vivo* to insulin. Effects of branched-chain amino acids. *Biochem J* 1988; 254: 579-584
 36. Tessari P, Trevisan R, Inchiostro S, Biolo G, Nosadini R, De Kreutzenberg SV, Duner E, Tiengo A, Crepaldi G: Dose-response curves of effects of insulin on leucine kinetics in humans. *Am J Physiol* 1986; 251: E334-E342
 37. Tessari P, Biolo G, Inchiostro S, Saggini L, Piccoli A, Tiengo A: Relationship between plasma leucine concentration and clearance in normal and type 1 diabetic subjects. *Acta Diabetol* 1992; 29: 6-10
 38. Hummel RP 3rd, Warner BW, James JH, Hasselgren PO, Fischer JE: Effects of indomethacin and leupeptin on muscle cathepsin B activity and protein degradation during sepsis. *J Surg Res* 1988; 45: 140-144
 39. Tessari P, Inchiostro S, Biolo G, Vincenti E, Sabadin L: Effects of acute systemic hyperinsulinemia on forearm muscle proteolysis in healthy man. *J Clin Invest* 1991; 88: 27-33
 40. Tessari P, Nissen SL, Miles JM, Haymond MW: Inverse relationship of leucine flux and oxidation to free fatty acid availability *in vivo*. *J Clin Invest* 1986; 77: 575-581
 41. Tessari P, Inchiostro S, Biolo G, Trevisan R, Fantin G, Marescotti MC, Iori E, Tiengo A, Crepaldi G: Differential effects of hyperinsulinemia and hyperaminoacidemia on leucine-carbon metabolism *in vivo*. Evidence for distinct mechanisms in regu-