

8. Bensinger WI, Weaver CH, Appelbaum FR, Rowley S, Demirer T, Sanders J, Storb R, Buckner CD: Transplantation of allogeneic peripheral blood stem cells mobilized by recombinant human granulocyte colony-stimulating factor. *Blood* 1995; 85: 1655–1658
9. Wagner JE, Kernan NA, Steinbuch M, Broxmeyer HE, Gluckman E: Allogeneic sibling umbilical-cord-blood transplantation in children with malignant and non-malignant diseases. *Lancet* 1995; 346: 214–219
10. Rubinstein P, Rosenfield RE, Adamson JW, Stevens CE: Stored placental blood for unrelated bone marrow reconstitution. *Blood* 1993; 81: 1679–1690
11. Gluckman E, Wagner J, Hows J, Kernan N, Bradley B, Broxmeyer HE: Cord blood banking for hematopoietic stem cell transplantation: an international cord blood transplant registry. *Bone Marrow Transplant* 1993; 11: 199–200
12. Catani L, Gugliotta L, Campanini E, Mangianti S, Gibellini D, Baravelli S, Vianelli N, Lemoli RM, Tura S: Megakaryocyte progenitors derived from bone marrow or G-CSF-mobilized peripheral blood CD34 cells show a distinct phenotype and responsiveness to interleukin-3 (IL-3) and PEG-recombinant human megakaryocyte growth and development factor (PEG-rHuMGDF). *Br J Haematol* 1998; 100: 207–218
13. Urashima M, Hoshi Y, Akiyama M, Kamijo M, Shishikura A, Kato Y, Akatsuka J, Maekawa K: Effects of cytokines on hematopoietic progenitor cells in cord blood, in bone marrow, and in peripheral blood mobilized by chemotherapy and G-CSF. *Acta Paediatr Jpn* 1995; 37: 21–26
14. Gehling UM, Ryder JW, Hogan CJ, Hami L, McNiece I, Franklin W, Williams S, Helm K, King J, Shpall EJ: *Ex vivo* expansion of megakaryocyte progenitors: effect of various growth factor combinations on CD34+ progenitor cells from bone marrow and G-CSF-mobilized peripheral blood. *Exp Hematol* 1997; 25: 1125–1139
15. Hao QL, Shah AJ, Thiemann FT, Smogorzewska EM, Crooks GM: A functional comparison of CD34+ CD38– cells in cord blood and bone marrow. *Blood* 1995; 86: 3745–3753
16. Van Epps DE, Bender J, Lee W, Schilling M, Smith A, Smith S, Unverzagt K, Law P, Burgess J: Harvesting, characterization, and culture of CD34+ cells from human bone marrow, peripheral blood, and cord blood. *Blood Cells* 1994; 20: 411–423
17. van de Ven C, Ishizawa L, Law P, Cairo MS: IL-11 in combination with SLF and G-CSF or GM-CSF significantly increases expansion of isolated CD34+ cell population from cord blood vs. adult bone marrow. *Exp Hematol* 1995; 23: 1289–1295
18. Liu J, Li K, Yuen PM, Fok TF, Yau FW, Yang M, Li CK: *Ex vivo* expansion of enriched CD34+ cells from neonatal blood in the presence of thrombopoietin, a comparison with cord blood and bone marrow. *Bone Marrow Transplant* 1999; 24: 247–252
19. Bender JG, Unverzagt K, Walker DE, Lee W, Smith S, Williams S, Van Epps DE: Phenotypic analysis and characterization of CD34+ cells from normal human bone marrow, cord blood, peripheral blood, and mobilized peripheral blood from patients undergoing autologous stem cell transplantation. *Clin Immunol Immunopathol* 1994; 70: 10–18
20. Steen R, Tjonnfjord GE, Egeland T: Comparison of the phenotype and clonogenicity of normal CD34+ cells from umbilical cord blood, granulocyte colony-stimulating factor-mobilized peripheral blood, and adult human bone marrow. *J Hematother* 1994; 3: 253–262
21. De Bruyn C, Delforge A, Bron D, Bernier M, Massy M, Ley P, de Hemptinne D, Stryckmans P: Comparison of the coexpression of CD38, CD33 and HLA-DR antigens on CD34+ purified cells from human cord blood and bone marrow. *Stem Cells* 1995; 13: 281–288
22. Drygalski A, Xu G, Constantinescu D, Kashiwakura I, Farley T, Dobrila L, Rubinstein P, Adamson JW: The frequency and proliferative potential of megakaryocytic colony-forming cells (Meg-CFC) in cord blood, cytokine-mobilized peripheral blood and bone marrow, and their correlation with total CFC numbers: implications for the quantitation of meg-CFC to predict platelet engraftment following cord blood transplantation. *Bone Marrow Transplant* 2000; 25: 1029–1034
23. Ratajczak J, Machalinski B, Samuel A, Pertusini E, Majka M, Czajka R, Ratajczak MZ: A novel serum free system for cloning human megakaryocytic progenitors (CFU-Meg). The role of thrombopoietin and other cytokines on bone marrow and cord blood CFU-Meg growth under serum free conditions. *Folia Histochem Cytobiol* 1998; 36: 61–66
24. Gordon MY, Lewis JL, Scott MA, Roberts IA, Goldman JM: Probability of progenitor renewal (PPR) and production of clonogenic progeny (CFU-GM) by primitive adherent progenitor cells in adult human bone marrow and umbilical cord blood. *Br J Haematol* 1995; 90: 744–749
25. Migliaccio G, Baiocchi M, Hamel N, Eddleman K, Migliaccio AR: Circulating progenitor cells in human ontogenesis: response to growth factors and replating potential. *J Hematother* 1996; 5: 161–170
26. Wyrsh A, dalle Carbonare V, Jansen W, Chklovskaja E, Nissen C, Surbek D, Holzgreve W, Tichelli A, Wodnar-Filipowicz A: Umbilical cord blood from preterm human fetuses is rich in committed and primitive hematopoietic progenitors with high proliferative and self-renewal capacity. *Exp Hematol* 1999; 27: 1338–1345
27. Migliaccio G, Migliaccio AR, Druzin ML, Giardina PJ, Zsebo KM, Adamson JW: Effects of recombinant human stem cell factor (SCF) on the growth of human progenitor cells in vitro. *J Cell Physiol* 1991; 148: 503–509
28. Migliaccio G, Migliaccio AR, Druzin ML, Giardina PJ, Zsebo KM, Adamson JW: Long-term generation of colony-forming cells in liquid culture of CD34+ cord blood cells in the presence of recombinant human stem cell factor. *Blood* 1992; 79: 2620–2627
29. Wang JC, Doedens M, Dick JE: Primitive human hematopoietic cells are enriched in cord blood compared with adult bone marrow or mobilized peripheral blood as measured by the quantitative in vivo SCID- repopulating cell assay. *Blood* 1997; 89: 3919–3924
30. Vaziri H, Dragowska W, Allsopp RC, Thomas TE, Harley CB, Lansdorp PM: Evidence for a mitotic clock in human hematopoietic stem cells: loss of telomeric DNA with age. *Proc Natl Acad Sci USA* 1994; 91: 9857–9860
31. Zandstra PW, Conneally E, Piret JM, Eaves CJ: Ontogeny-associated changes in the cytokine responses of primitive human hematopoietic cells. *Br J Haematol* 1998; 101: 770–778
32. Lansdorp PM, Dragowska W, Mayani H: Ontogeny-related changes in proliferative potential of human hematopoietic cells. *J Exp Med* 1993; 178: 787–791
33. Deutsch VR, Olson TA, Nagler A, Slavin S, Levine RF, Eldor A: The response of cord blood megakaryocyte progenitors to IL-3, IL-6 and aplastic canine serum varies with gestational age. *Br J Haematol* 1995; 89: 8–16
34. Weckx SF, Van Bockstaele DR, Plum J, Moulijn A, Rodrigus I, Lardon F, De Smedt M, Nijs G, Lenjou M, Loquet P, Berneman ZN, Snoeck HW: CD34++ CD38– and CD34+ CD38+ human hematopoietic progenitors from fetal liver, cord blood, and adult bone marrow respond differently to hematopoietic cytokines depending on the ontogenic source. *Exp Hematol* 1998; 26: 1034–1042
35. Nishihira H, Toyoda Y, Miyazaki H, Kigasawa H, Ohsaki E: Growth of macroscopic human megakaryocyte colonies from cord blood in culture with recombinant human thrombopoietin